

Review

Invasive smallmouth bass (*Micropterus dolomieu*): history, impacts, and control

Grace L. Loppnow^{1*}, Kris Vascotto² and Paul A. Venturelli¹

¹ 135 Skok Hall, 2003 Upper Buford Circle, St. Paul, MN 55108, USA

² Ontario Ministry of Natural Resources, 190 Cherry Street, Chapleau, ON P0M 1K0, Canada

E-mail: lopp0010@umn.edu (GLL), kvascotto@gmail.com (KV), pventure@umn.edu (PAV)

*Corresponding author

Received: 16 February 2013 / Accepted: 30 July 2013 / Published online: 2 September 2013

Handling editor: Kathleen Beyer

Abstract

In this review, we (i) describe smallmouth bass (*Micropterus dolomieu* Lacepède, 1802) invasions past, present, and future; (ii) summarize the impact that this species can have on native communities; and (iii) describe and discuss various options for control. *M. dolomieu* are invasive throughout much of the United States, southern portions of Canada, and in countries in Europe, Asia, and Africa. Historically, this species spread via stocking programs intended to improve sport fisheries. Currently, their spread is facilitated by anglers and global climate change. Models predict that *M. dolomieu* will continue to spread with consequences for native prey fish, sport fish, and food webs through predation, competition, and hybridization. Effective control methods are necessary to mitigate these impacts. Options for *M. dolomieu* control include biological control, chemical control, environmental manipulation, and physical removal. However, our review of the literature suggests that only a handful of the possible control options have been explored (usually in isolation and with limited success), and that there is a clear need for focused research and informed management. For example, our elasticity analysis of published *M. dolomieu* matrix population models suggests that *M. dolomieu* control will be most effective when it targets eggs, larvae, and juveniles. We recommend targeting these life stages by using nest failure as part of an adaptive and integrated pest management approaches that incorporate existing and emerging technologies. However, we also emphasize that *M. dolomieu* control, where necessary and possible, is more likely to take the form of suppression rather than permanent eradication. Therefore, we also recommend efforts to prevent *M. dolomieu* (re)introduction.

Key words: largemouth bass; black bass; invasive species; fisheries management; integrated pest management; climate change

Introduction

The smallmouth bass (*Micropterus dolomieu* Lacepède, 1802) is a cool-warm water centrarchid (Brown et al. 2009; Shuter et al. 1980, 1989) and a popular sport fish among anglers. *M. dolomieu* are littoral predators, generally consuming small prey fish and crayfish (Vander Zanden et al. 1999). During the spring, male *M. dolomieu* build and guard nests in the shallows of lakes and streams (Ridgway et al. 1991). *M. dolomieu* are native to freshwater systems in 23 states in the east-central United States (Rahel 2000; Fuller and Cannister 2011) and the southern portions of two Canadian provinces (Scott and Crossman 1973).

Outside of its native range, *M. dolomieu* is an invasive species for which there is currently no

effective means of control. *M. dolomieu* are invasive across much of the United States, southern portions of Canada, and in 9 other countries throughout the world (Fuller and Cannister 2011; Lyons 2011; Iguchi et al. 2004a). As with many invasive fishes, the spread of *M. dolomieu* beyond its native range has been facilitated by intentional and accidental stocking and climate-mediated habitat expansion (Jackson 2002; Rahel and Olden 2008). Invasive *M. dolomieu* reduce native small-bodied fish abundance and diversity through predation, outcompete other piscivorous game fish, and indirectly change planktonic and benthic communities (Jackson 2002). To date, most attempts to control invasive *M. dolomieu* have either produced undesirable results (e.g., Zipkin et al. 2008) or proven prohibitively labor-intensive (e.g., Tyus and Saunders 2000).

Here, we review the literature on invasive *M. dolomieu* and consider various options for control or eradication. Where *M. dolomieu* literature is lacking, we draw from the literature on invasive largemouth bass (*Micropterus salmoides* Lacepède, 1802), a sister species with a similar life history. First, we summarize the history of *M. dolomieu* invasions and review research that predicts *M. dolomieu* spread with climate change. We then summarize the impacts that invasive *M. dolomieu* have on native species and food webs. Finally, we outline various options for controlling or eradicating invasive species and describe their relevance to *M. dolomieu* in light of life history and previous control attempts.

Smallmouth bass invasions: Past, present, and future

Movement of *M. dolomieu* beyond their native range began primarily through intentional stocking by fisheries managers during the 19th century. The prevailing attitude in fisheries and wildlife management during this period was that nature should be controlled and improved upon. Additionally, recreational fishing was just beginning to come into its own during this era. Books aimed at outdoorsmen promoted bass fishing (e.g. Henshall 1889 and 1903) which was becoming a popular sport. Consequently, managers believed that introducing *M. dolomieu* would prove beneficial. The building enthusiasm for bass fishing led to a wealth of research on bass spawning and rearing (Bower 1897; Cushman 1917; Lydell 1902; Ripple 1908) allowing hatcheries to produce *M. dolomieu* for stocking throughout North America. For example, *M. dolomieu* were introduced to California in 1874 to improve sport fisheries (Moyle 1976). Similarly, from 1868 to 1881 the Maine Commissioners of Fisheries not only authorized *M. dolomieu* introductions in 51 water bodies, but also encouraged indiscriminate introduction by the public (Warner 2005). In the early 20th century, managers introduced *M. dolomieu* into lakes in Ontario, Alberta and Manitoba, and even into a national park in Saskatchewan (Rawson 1945). Unfortunately, fisheries managers in the 19th and early 20th century knew little about both the importance of native fishes and the threats that non-native fishes such as *M. dolomieu* could pose to biodiversity. Indeed, non-native fishes were often introduced to control certain “undesirable” native species and enrich biodiversity (Hey 1926; Moyle

1976). Some jurisdictions even enacted laws to protect these non-natives (Cambray 2003; Hey 1926). Enthusiasm for non-native stocking was eventually tempered by changing attitudes and a better understanding of the impacts of non-native fishes. Stocking non-native fish to provide angling opportunities lost momentum after a final surge in the 1950s (Crossman 1991) and by the 1980s most authorized introductions were into systems that had already been invaded (e.g. Rahel 2004; Carey et al. 2011).

Intentional and unintentional introductions by anglers have been and continue to be major drivers behind the spread of *M. dolomieu* (Jackson 2002). Unintentional introductions of non-native fishes commonly result from bait bucket transfers (Litvak and Madrak 1993). The most comprehensive analysis to date (Drake 2011) suggests that bait bucket transfers in Ontario are responsible for as many as 20 introduction events of *M. dolomieu* per system per year into waterbodies outside of their current range. Intentional introductions occur because *M. dolomieu* are a popular sport fish. Bass are responsible for millions of angler fishing days per year in the Pacific Northwest (Carey et al. 2011), and 77.8% of competitive fishing events in North America’s inland waters (Schramm et al. 1991). Both casual and competitive angling are important sources of revenue and development for many communities (Chen et al. 2003). Bass fishing has such a positive image that the negative effects of bass introduction usually go ignored. As early as the late 19th century, citizens recognized that introduced *M. dolomieu* alter fish assemblages (Warner 2005), but the general public has remained apathetic towards the spread of *M. dolomieu* (Jackson 2002). Anglers continue to intentionally introduce *M. dolomieu* to create more fishing opportunities for bass, without knowing or acknowledging the potential impacts.

Recently, the spread and establishment of *M. dolomieu* has also been facilitated by global climate change. The establishment of *M. dolomieu* is dependent on temperature because their range is limited by the severity of overwintering stress in coldwater lakes (Shuter et al. 1980; Shuter et al. 1989; Jackson et al. 2001). Suitable habitat for *M. dolomieu* is expanding because of warming of lakes and streams attributed to global climate change. Climate change can also facilitate the spread of *M. dolomieu* to uninvaded systems through flooding associated with an increase in extreme weather events (Rahel and Olden 2008).




Figure 1. The invasion status of smallmouth bass worldwide. Data are from Table 1, where “introduced” has been interpreted to mean that an invasive population has established. See Fuller and Cannister 2011 and Brown et al. 2009 for details on native and invasive ranges within the United States and Canada, respectively.

Flooding has contributed to the spread of at least two other species: bighead carp (*Hypophthalmichthys nobilis* Richardson, 1845) and silver carp (*Hypophthalmichthys molitrix* Valenciennes, 1844) (Kolar et al. 2005). *M. dolomieu* are now spreading northward into Canada, where they pose a serious threat to native species (Dextrase and Mandrak 2006). At current temperatures, 6% of Ontario lakes are predicted to be at high risk for *M. dolomieu* introduction, establishment, and subsequent impacts on native fauna (Vander Zanden et al. 2004); with climate change this number could increase to 20% by the year 2100 (Sharma et al. 2009b). A conservative estimate is that at least 50% of Canada will become thermally suitable *M. dolomieu* habitat, including some arctic locations (Sharma et al. 2007). Combining these predicted changes in habitat suitability with predictions of where introductions will occur suggests that Manitoba and Ontario are the provinces at greatest risk for *M. dolomieu* invasion (Chu et al. 2005). Similar increases in thermal habitat are expected at northern latitudes throughout the world. For example, although cold temperatures prevented the establishment of *M. dolomieu* introduced to Sweden in the 1960s (Curry-Lindahl 1966, Kullander et al. 2012), they may be able to establish there under a warmer climate.

As a result of introductions and habitat expansion, *M. dolomieu* are currently invasive throughout

much of the United States, southern portions of Canada, Mexico, and 8 countries on three other continents (Figure 1, Table 1). Forty-two states (including Hawaii) and most Canadian provinces bordering the United States have *M. dolomieu* in areas where they are considered non-native (Fuller and Cannister 2011; Lyons 2011). *M. dolomieu* have also established viable populations in Europe, Africa, and most notably in Japan where habitat suitability models suggest that most waterbodies in the country are at risk of *M. dolomieu* invasion (Iguchi et al. 2004b).

Impacts of invasive smallmouth bass

Invasive *M. dolomieu* can disrupt the native ecology of the systems to which they have been introduced. Bass are voracious predators that can decrease the abundance of, change the habitat used by, and even extirpate small prey fish such as brook stickleback (*Culaea inconstans* Kirtland, 1840), fathead minnow (*Pimephales promelas* Rafinesque, 1820), pearl dace (*Margariscus margarita* Cope, 1867), finescale dace (*Phoxinus neogaeus* Cope, 1867) and northern redbelly dace (*Phoxinus eos* Cope, 1861) (MacRae and Jackson 2001; Trumpickas et al. 2011). In streams, prey fish alter their behavior to avoid invasive *M. dolomieu* by moving from pools to riffles and areas with more structural complexity

Table 1. Known smallmouth bass introductions by country. Data compiled from sources listed as well as FishBase and the FAO Fisheries and Aquaculture Department. Table does not include Belize because this record is believed to be in error (Peter Esselman, pers. comm.).

Country	Status	Year introduced	Introduced from (country)	Method of introduction	Sources
Austria	introduced	unknown	unknown	unknown	Welcomme 1988
Belgium	introduced	1873	USA	angling/sport	Welcomme 1988
Canada	native and introduced	unknown	USA, Canada	angling/sport	Scott and Crossman 1973
Czech Republic	introduced	1889	unknown	unknown	Hanel 2003
Denmark	not established	1958	Canada	unknown	Ostergaard pers. comm.
Fiji	not established	1962	unknown	angling/sport	Andrews 1985
Finland	not established	1893-1963	Sweden, Germany, Canada, Sweden, Germany, Canada	angling/sport, aquaculture	FAO 1997
France	not established	unknown	North America	unknown	Allardi and Keith 1991
Germany	not established	1880	USA	angling/sport	Welcomme 1988
Guam	not established	1962	unknown	unknown	Welcomme 1988
Japan	introduced	unknown	unknown	unknown	Masuda et al. 1984
Mauritius	introduced	unknown	unknown	unknown	Fricke 1999
Mexico	introduced	1975	USA	aquaculture	Welcomme 1988
Netherlands	not established	1984	USA	unknown	Welcomme 1988
Norway	not established	1887-1895	Germany	fill ecological niche	Welcomme 1988
Slovakia	introduced	unknown	unknown	unknown	Welcomme 1988
South Africa	introduced	1937	USA	angling/sport	Welcomme 1988
Swaziland	not established	1938	South Africa	angling/sport	Welcomme 1988
Sweden	not established	1890	USA, Germany	angling/sport	Welcomme 1988
Tanzania	introduced	unknown	unknown	unknown	Fermon 1997
United Kingdom	not established	1878-1890	USA	angling/sport	Welcomme 1988
United States	native and introduced	unknown	USA	angling/sport	Page and Burr 1991
Vietnam	introduced	unknown	South Africa	unknown	Kuronuma 1961
Zimbabwe	not established	1942	South Africa	angling/sport	Welcomme 1988

(Schlosser 1987). Shifting habitat use from pools to shallower areas could expose these fish to predation from terrestrial predators and result in higher energy expenditures during foraging. Invasive *M. dolomieu* pose a serious predatory threat to small native fish in the Yampa River, Colorado, a regional hotspot of native fish diversity (Johnson et al. 2008). The rate of piscivory by *M. dolomieu* is estimated to be ten times that of two other invasive piscivores in this system. In New Mexico, predation by invasive *M. dolomieu* is depleting populations of the threatened bigscale logperch (*Percina macrolepida* Stevenson, 1971) (Archdeacon and Davenport 2010). The spread of *M. dolomieu* into Ontario alone is expected to extirpate more than 25,000 cyprinid populations (Jackson and Mandrak 2002). The loss of such species can lead to both a loss of diversity within invaded waters and a homogenization of fish fauna among invaded waters (MacRae and Jackson 2001; Jackson 2002).

Invasive *M. dolomieu* can also impact top predators, many of which are also prized sport fish. These impacts occur primarily through competition for prey and predation on juveniles. Salmon and trout are particularly sensitive to *M. dolomieu* invasion (Sharma et al. 2009a). Stable isotope studies suggest that *M. dolomieu* predation alters food webs by forcing piscivorous lake trout (*Salvelinus namaycush* Walbaum in Artedi, 1792) to prey on zooplankton, a low-quality food source (Vander Zanden et al. 1999; Morbey et al. 2007). However, the presence of pelagic prey fish can buffer *S. namaycush* from the effects of competition with *M. dolomieu* by providing an alternative high - quality food source (Vander Zanden et al. 2004). In the event of a shift to sub-optimal prey, *S. namaycush* growth and reproduction could be limited. For example, in the 1960s *M. dolomieu* were introduced into Utah's Flaming Gorge Reservoir to control the native Utah chub (*Gila atraria* Girard, 1856),

Table 2. Invasive fish control methods, examples of their use for smallmouth bass control, the bass life stage they target, and their pros and cons. This table draws from and builds upon a review by Halfyard (2010). For “Target”, 1=eggs, 2=fry, 3=juveniles, 4=adults.

Method	Description	Example(s)	Target	Pros	Cons
Biological control (pathogens)	Introduction of a parasite or disease that targets bass	Davis (1937,1942)*, McCormick and Stokes (1982)*, Grizzle et al. 2003*	1-4	inexpensive (application), not labor-intensive, effective in all waterbodies and habitats	expensive (development), unconventional, controversial, risk to non-target species, resistance
Biological control (predators)	Introduction of organisms that prey on young bass	Iguchi and Yodo (2004)*	1,2	inexpensive, not labor-intensive, effective in all waterbodies and habitats	controversial, unexpected ecological effects
Biological control (sterilization)	Limit reproductive success (e.g. sterile males)	Dey et al. (2010)*	1,4	species-specific, effective in all waterbodies and habitats	expensive, labor-intensive, unconventional
Chemical	Use of piscicides to kill bass	Smith (1941), Ward (2005)	1-4	not labor-intensive, effective in all waterbodies and habitats	unconventional, controversial, expensive, affects non-target species, destructive
Environmental manipulation (water level)	Complete or partial dewatering to affect survival/reproduction	Kleinschmidt (2008), Mukai et al. 2011*, Kitazima and Mori 2011*	1-4	effective, inexpensive, not labor-intensive	affects non-target species, controversial, limited applicability
Environmental manipulation (winterkill)	Encouragement of a low-oxygen environment that cannot support bass	Smale and Rabeni 1995*, Verrill and Berry 1995*, Shroyer 2007*	3,4	effective, inexpensive, not labor-intensive	unconventional, affects non-target species, limited applicability
Removal (angling)	Use of angling to remove bass	Boucher (2006)	3,4	conventional, uncontroversial, species- and size-selective, applicable to all depths	labor-intensive, inefficient, impractical in large waterbodies
Removal (electrofishing)	Use of electrofishing gear to remove bass	Rinne (2001), Weidel et al. (2007), Boucher (2006, 2005), Burdick (2008), Hawkins et al. (2008)	3, 4	conventional, uncontroversial, effective in small waterbodies	labor-intensive, inefficient, affects non-target species, ineffective in deep/complex habitat or large waterbodies, overcompensation
Removal (explosives)	Use of explosives to kill bass	Munther (1970)*, Metzger and Shafland 1986*	2-4	cheap and effective in small waterbodies and in all habitats	unconventional, controversial, affects non-target species, destructive, dangerous, ineffective in large waterbodies
Removal (netting)	Use of nets and traps to remove bass	Boucher (2006), Gomez and Wilkinson (2008)	3,4	conventional, uncontroversial, species- and size-selective, applicable to all depths	labor-intensive, ineffective, affects non-target species

*Study contains proof of concept of the applicability of the control method to smallmouth bass, but does not attempt smallmouth bass control.

which was competing with salmonid sport fish (Teuscher and Luecke 1996). However, decades after introduction, competition with *M. dolomieu* for food appears to be inhibiting the growth of young *S. namaycush* in the reservoir (Yule and Luecke 1993). In Canada, climate change models predict that by 2100, 11% of *S. namaycush* populations will be negatively impacted by competition with *M. dolomieu* (Sharma et al. 2009b).

Invasive *M. dolomieu* also impact sport fish by preying directly on juveniles. For example, predation by *M. dolomieu* is putting some threatened and endangered species of Pacific salmon (*Oncorhynchus* spp. Suckley, 1861) at

greater risk of extinction (Reiman et al. 1991; Carey et al. 2011). In the Pacific Northwest, invasive *M. dolomieu* consume an average of about 20% of outmigrating juvenile salmon in streams; in some cases that figure can approach 40% (Sanderson et al. 2009). *M. dolomieu* also consume young walleye (*Sander vitreus* Mitchell, 1818) (Liao et al. 2004), but the extent to which this predation affects *S. vitreus* populations is unclear. On one hand, the native range of *S. vitreus* includes the native range of *M. dolomieu* and these species appear capable of coexisting in both native (e.g., Johnson and Hale 1977; Kempinger and Carline 1977) and non-native

(Galster et al. 2012) *M. dolomieu* lakes. On the other hand, population abundances can be inversely related, likely due to multiple factors including predation (Fayram et al. 2005; Johnson and Hale 1977) and preferences for different conditions (Inskip and Magnuson 1983; Robillard and Fox 2006).

Invasive *M. dolomieu* can also hybridize with native bass. Hybridization can result in genetic introgression and the displacement, decline or extirpation of native species. *M. dolomieu* are known to hybridize with *M. salmoides*, spotted bass (*Micropterus punctulatus* Rafinesque, 1819), and Guadalupe bass (*Micropterus treculii* Vaillant and Bocourt, 1874) (Whitmore 1983; Whitmore and Hellier 1988). Hybridization with *M. treculii* is of particular concern because this species is endemic to the Edwards Plateau of south central Texas (Edwards 1980).

Fish are not the only taxa affected by invasive *M. dolomieu*; mammals, birds, amphibians, reptiles, and invertebrates can be impacted as well. *M. dolomieu* will consume almost any prey small enough to ingest including crayfish, rats, mice, young waterfowl, frogs, snakes, and salamanders (Sanderson et al. 2009). Frog species can be impacted by predation from *M. dolomieu*, although the severity could depend on the presence of other invasive species and the life stage of the frog (Kiesecker and Blaustein 1998). Any organism that depends on the prey of *M. dolomieu* can also be impacted. In an extreme case, competition with invasive *M. dolomieu* and *M. salmoides* for food contributed to the extinction of an endemic Guatemalan waterbird, the Atilán grebe (*Podilymbus gigas* Griscom, 1929) (Hunter 1988). In ponds invaded by *M. salmoides*, the loss of prey fish and crayfish populations can lead to a reduction in top-down control of benthic invertebrates and macrophytes (Maezono and Miyashita 2003; Maezono et al. 2005).

Controlling invasive smallmouth bass

M. dolomieu control is essential for mitigating, minimizing and perhaps even eliminating the impacts of *M. dolomieu* on native species and food webs. However, there has been little documented work on *M. dolomieu* control to date, and many potential control options remain untested. Here we describe and discuss various control options as they relate to *M. dolomieu*. This section draws from and builds upon a recent review by Halfyard (2010). We summarize control options in Table 2.

Removal

Removal refers to the physical capture and removal of fish from a system, typically via electrofishing, netting, explosives, or angling. These methods are labor-intensive and rarely result in successful control. However, in certain systems they may be an effective component of an integrated management plan.

Electrofishing

Electrofishing is a common, uncontroversial removal method in fisheries management that has been applied to invasive *M. dolomieu* with limited success. Electrofishing programs in small reaches of the Colorado and Yampa Rivers decreased *M. dolomieu* abundance, but only temporarily due to immigration (Burdick 2008; Hawkins et al. 2008). Other attempts to control invasive *M. dolomieu* via electrofishing have ultimately failed because of increased recruitment following treatment (Boucher 2005 and 2006; Weidel et al. 2007; Hawkins et al. 2008). In one striking example, the mass removal of 47,474 *M. dolomieu* over a 6-year period from an Adirondack lake in New York initially reduced *M. dolomieu* abundance by 90%, but ultimately resulted in increased abundance (Weidel et al. 2007; Zipkin et al. 2008). This unexpected increase was attributed to decreased intraspecific competition that led to accelerated maturation of juveniles and, ultimately, improved recruitment (Ridgway et al. 2002; Zipkin et al. 2008). This phenomenon is known as the hydra effect or overcompensation (Abrams 2009; Strevens and Bonsall 2011; Zipkin et al. 2008). Because electrofishing gear tends to remove more adults than juveniles (Moore et al. 1986; Kulp and Moore 2000; Earle and Lajeunesse 2007), this method can lead to overcompensation. Therefore, a control plan that involves electrofishing should include one or more methods that reduce the abundance of young *M. dolomieu*. Electrofishing is only likely to be effective in shallow, isolated streams and ponds absent of complex habitat. This method is labor-intensive, inefficient, non-species-specific, and requires repeated, long-term application.

Netting

Netting is another common fisheries management tool but is generally ineffective at controlling invasive *M. dolomieu*. For example, a springtime

attempt to net invasive *M. dolomieu* in a 70 ha pond captured only 7 individuals in 2,103 trap net hours (0.003 fish/hour), as compared to 200 *M. dolomieu* captured in 8.62 hours of electrofishing effort (23.2 fish/hour; Boucher 2006). In another attempt, less than 1% of the 3083 fish captured in gill nets were *M. dolomieu* (Gomez and Wilkinson 2008). These results might reflect the relative abundance of *M. dolomieu* in the system, the low vulnerability of centrarchids to passive netting (Hayes et al. 1996), or seasonal variation in catchability (i.e., *M. dolomieu* are most trappable in mid-summer, Wright 2000). In general, nets are much less effective at catching *M. dolomieu* than electrofishing (Bacula et al. 2011 and references therein). Therefore, although netting is a familiar, available, and uncontroversial control option that can be both size- and species-selective and deployed at most depths, the inability of nets to catch large numbers of *M. dolomieu*, even when effort is high, is a significant shortcoming that precludes the use of nets for control, even in combination with other methods. However, netting may be the only option in systems that are not conducive to electrofishing. For example, invasive young-of-the-year *M. dolomieu* in Lake Opeongo, a low-conductivity lake in Algonquin Park, Canada, are routinely (and effectively) sampled using minnow traps (e.g. Dunlop et al. 2005a, 2005b).

Explosives

Explosives have been proposed for invasive fish control (Lee 2001). Although there are no known applications to *M. dolomieu* control, the use of detonation cord to sample *M. dolomieu* in deep reaches of the Middle Snake River, Idaho (Munther 1970) suggests that at least some degree of removal is possible. Detonation cord tends to kill most fish within 9 meters of the blast (Metzger and Shafland 1986). In general, explosives are effective at killing adult and larval fish with swim bladders, but not fish eggs (Baxter II et al. 1982; Metzger and Shafland 1986; Bayley and Austen 1988; Keevin et al. 2002; Settle et al. 2002; Faulkner et al. 2008).

Explosives are an effective and relatively cheap method for killing fish in almost any habitat, but there are a number of issues that are likely to limit their use in *M. dolomieu* control. Explosives are difficult to obtain, highly controversial, and dangerous to both human and environmental health. They are not selective for

fish species, and can destroy habitat and leave behind toxic chemical residues (Hayes et al. 1996; Lotufo and Lydy 2005). Explosives may also be inappropriate for large waterbodies because of scale. Several authors recommend explosives only when there are no other options for sampling or control (e.g., Bayley and Austen 1988; Hayes et al. 1996).

Angling

Removal by angling is another control option that is unlikely to be effective. In the only documented attempt, anglers removed *M. dolomieu* from a 70 ha pond in Maine at a rate of 0.31 fish/hour as compared to 23.2 fish/hour for electrofishing (Boucher 2006). The author did not believe that angling was an appropriate control measure for that system. In another program, nearly 300 angler hours over two years resulted in the removal of just 150 *M. dolomieu* (Gomez and Wilkinson 2008). Even though angling for *M. dolomieu* control has not been successful, increased fishing pressure either from liberalized regulations or intensive effort has been advocated for the control of other invasive fishes (Wydoski and Wiley 1999; Beamesderfer et al. 1996; Moore et al. 2005). Because angling is an inefficient removal method, it is probably most effective in small systems and/or when fishing pressure is high.

Options for enhancing removal

The efficacy of a particular removal method can be enhanced through techniques that improve catchability. Here we describe two such techniques: pheromone-baited traps and “Judas fish”. Pheromone-baited traps use species-specific chemical attractants to improve trap efficiency (Sorensen and Stacey 2004). Although we are unaware of examples involving *M. dolomieu*, these traps have been used successfully for other invasive fish such as sea lamprey (*Petromyzon marinus* Linnaeus, 1758) and common carp (*Cyprinus carpio* Linnaeus, 1758) (Wagner et al. 2006; Sorensen and Stacey 2004). Therefore, we recommend exploring pheromone traps as an option for *M. dolomieu* control. Another technique that can improve catchability is the addition of “Judas fish” to the system (Bajer et al. 2011). The Judas technique was first developed in Hawaii, where it was used to locate non-native feral goats (Taylor and Katahira 1988). A Judas

fish is a conspecific that has been implanted with a radio tag and then tracked to an aggregation of fish. These fish can then be more effectively targeted for removal. The Judas fish technique should be evaluated for its potential to locate aggregations of *M. dolomieu* (e.g., during spawning or winter shoaling), particularly in large systems.

Chemical

The addition of chemical piscicides to a system is perhaps the most common method of fish control (Wydoski and Wiley 1999; Meronek et al. 1996). Piscicides such as rotenone and antimycin-A are effective at killing a large proportion of fish in the system with minimal effort (Lennon et al. 1971, Baker et al. 2008). Both rotenone and a newly-developed piscicide, Supaverm®, can effectively kill invasive *M. dolomieu* (Smith 1941; Ward 2005). Rotenone eliminated *M. dolomieu* from Potter's Lake, New Brunswick (46 ha), and Supaverm® holds promise as a more selective option that tends to spare native minnows. Overall, and given adequate funding, chemical control can be a quick and effective option for *M. dolomieu* control in many systems. However, most piscicides are lethal to all fish in the system (Finlayson 2001; Dawson and Kolar 2003) and can also affect amphibians, aquatic invertebrates, and zooplankton (Smith 1940; Brown and Ball 1943; Morrison 1979; Finlayson et al. 2000; Arnekleiv et al. 2001; Ling 2001; Dinger and Marks 2007). Additionally, stakeholders may be opposed to this option on ethical grounds or due to its non-target effects and high cost. Managers should also be aware that piscicides are not always 100% effective at extirpating invasive fishes. Chemical eradication is, on average, only 35% effective 10 years after treatment (Wydoski and Wiley 1999). For chemical treatment to be effective, managers must acknowledge and manage for public opposition, non-target effects, and the potential need for additional control measures.

Biological control

Biological control (biocontrol) refers to the introduction or enhancement of an invasive species' predators or pathogens, or the sterilization of the invasive species. Although these methods tend to be controversial and are largely untested

for *M. dolomieu*, there is evidence to suggest that they can be effective. To this end, we encourage research into the efficacy of biocontrol methods as they apply to invasive *M. dolomieu*.

Predation

Predation is the most common and effective method of biological control, through either the introduction of new predators or the enhancement of existing ones (Wydoski and Wiley 1999). This type of control would be most effective for small non-game fish or young sport fish (e.g., *M. dolomieu* eggs, fry, and juveniles) given their vulnerability to predation. The impact of this kind of biocontrol on *M. dolomieu* may be limited by the tendency of nesting males to aggressively guard eggs and larvae from predators during the spawning season (Ridgway et al. 1991). However, nest guarding may not be a limiting factor in all cases. For example, native Japanese dace (*Tribolodon hakonensis* Günther, 1877) are extremely effective nest predators, consuming on average 92.4% of invasive *M. dolomieu* eggs, even while males guard their nests (Iguchi and Yodo 2004). In this case and others, removing the guarding male would enhance nest predation. Nest predators such as crayfish, yellow perch (*Perca flavescens* Mitchell, 1814), sunfish (*Lepomis* spp. Rafinesque, 1819), and the introduced round goby (*Neogobius melanostomus* Pallas, 1814) can consume an entire nest of unprotected eggs in as little as 17 minutes (Kieffer et al. 1995; Steinhart et al. 2004).

Although invasive species are unlikely to be appropriate predators to introduce for biocontrol, management plans that enhance native predators or introduce otherwise benign predators could be useful. Once an appropriate predator is identified, this type of control is usually inexpensive, requires minimal effort, and can be effective in a variety of habitats. Predatory biocontrol is nonetheless risky and controversial (Hoddle 2004). Before introducing a predator species (or enhancing natural predators), it is important to consider the vulnerability of the target species, the potential for non-target effects, and the likelihood that introduced predators will survive, establish, and spread (Wydoski and Wiley 1999). Managers should carefully weigh the pros and cons of this untested method and researchers should consider its study, as it appears to be a promising option for *M. dolomieu* control.

Pathogens and parasites

The introduction or enhancement of novel or existing pathogens or parasites is another biocontrol option for invasive fish. We find one promising example in Australia, where research is underway to use koi herpes virus to control common carp (McCull et al. 2007). Although we are unaware of attempts to use pathogens or parasites to control invasive *M. dolomieu*, there are at least two *M. dolomieu*-specific parasites that hold promise. The first parasite is a protozoan that attaches itself to *M. dolomieu* gills and can cause mortality (Davis 1937, 1942). The second parasite is a tapeworm (*Proteocephalus ambloplitis* Leidy) that limits *M. dolomieu* fecundity by preferentially infesting its oocytes (McCormick and Stokes 1982). With proper testing and development, these parasites could be used as biocontrol agents for invasive *M. dolomieu*. Similarly, the pathogen known as the largemouth bass virus (Family Iridoviridae; genus unknown) causes a disease that is specific to *M. salmoides* (Grizzle et al. 2003) but could potentially be genetically engineered to target *M. dolomieu*. Testing and developing *M. dolomieu*-specific pathogens and parasites is likely to be both expensive and time-consuming. Once developed, however, these agents could be used simply and cheaply in almost any system. Nonetheless, development and application should be undertaken with caution to avoid effects on non-target species, the development of resistance, and public controversy (Wydoski and Wiley 1999).

Sterilization

Sterilization is a form of biocontrol that involves the release of sterile conspecifics or the alteration of individual physiology to limit reproductive success. Sterilization is not currently available as a control technique for invasive *M. dolomieu*, but the idea merits further research. One promising option is to use pheromones to effectively sterilize *M. dolomieu* by altering their behavior. Treatment with pheromones can reduce a male bass's ability to guard his nest, causing increased nest failure (Dey et al. 2010). Another option is the release of sterile males, which has been used successfully for sea lamprey (Twohey et al. 2003). Recombinant gene therapy could be used to develop sterile *M. dolomieu*, but this

technology is largely untested (Thresher 2008). Other sterilization techniques such as irradiation and chemically-induced sterilization should be researched for *M. dolomieu*. Sterilization is species-specific and applicable to all types of habitats, but it can be expensive to develop, controversial, and labor-intensive to implement.

Environmental manipulation

Environmental manipulation seeks to control invasive fishes via changes to their physical environment. Here we focus on manipulation of water levels and dissolved oxygen concentration.

Water level manipulation

Draining, or dewatering, a system is considered to be the most effective way to guarantee complete removal of fish (Finlayson et al. 2002; Ling 2001; McClay 2000). However, this approach affects non-target species and organisms and is impractical in many places due to logistics or public opposition. Partial dewatering may be a more feasible option, especially in managed reservoirs, rivers, and streams in which it is easy to manipulate water levels. Partial dewatering can cause behavioral changes in *M. dolomieu* and increase predation on juveniles (Rogers and Bergersen 1995; Heman et al. 1969). Because *M. dolomieu* spawn in shallow water (Ridgway et al. 1991), partial dewatering can also be used to limit the availability of spawning habitat or kill incubating eggs. The only example of water level manipulation specifically targeted at controlling invasive *M. dolomieu* induced discharge pulses in a stream during the spawning season (Kleinschmidt 2008). These pulses evacuated all fry from 43% of the study nests and removed some of the fry from another 21%. While the author believes that this reduction in *M. dolomieu* young could improve trout habitat, the long-term impacts of this treatment have not been determined as of 2013. Water level management is probably an inexpensive and effective option for *M. dolomieu* control in certain systems. However, this method cannot be applied everywhere, has the potential for non-target effects, and could be controversial given conflicting water uses (agriculture, industry, residential, recreational, hydroelectric power, navigation, etc.).




Figure 2. Elasticities of the annual survival probability (s_i) and fertility (f_i) at age i for four smallmouth bass matrix population models.

Dissolved oxygen concentration

It may be possible to control *M. dolomieu* by reducing the wintertime concentration of dissolved oxygen (DO) below the lethal threshold of ~1.2 mg/L (Smale and Rabeni 1995). One can induce lethal concentrations of DO in winter (producing a winterkill event) by partially drawing down water or causing disturbances (e.g., artificial mixing of the water column) that stir up anoxic water and sediments containing oxygen-consuming bacteria (Verrill and Berry 1995; Shroyer 2007). Winterkill has been used successfully for other invasive fishes (Verrill and Berry 1995; Shroyer 2007), but not for *M. dolomieu*. The applicability of this option is likely limited to shallow, eutrophic lakes and ponds, where it could be effective and inexpensive. However, inducing low DO is likely to kill non-target organisms and could be controversial.

Targeting specific life stages

Directed research is a necessary first step in the development of *M. dolomieu* control methods that are both efficient and effective (Buhle et al. 2005; Eiswerth and Johnson 2002). First and foremost, research should help to focus management

effort by identifying those aspects of the *M. dolomieu* life cycle that contribute most to population growth rate. To this end, we conducted an elasticity analysis of four published matrix population models of *M. dolomieu* from lakes and rivers in Canada and the United States (Peterson and Kwak 1999; Van Winkle et al. 2001; Rose 2005; Marcinkevage-Miller 2007). Elasticity is a measure of the sensitivity of population growth rate to systematic and proportional changes to age-specific parameters for survival and fertility. Our results suggest that population growth rate is most sensitive to survival in the first 1–4 years of life (Figure 2). Therefore, from a biological perspective, targeting *M. dolomieu* eggs, larvae, and juveniles is the most efficient approach to controlling invasive *M. dolomieu*. Additionally, targeting these early life stages alone or in concert with management of adult *M. dolomieu* could potentially prevent overcompensation.

Options for targeting *M. dolomieu* eggs and larvae outnumber options for targeting *M. dolomieu* juveniles, and would be far more effective due to the relatively narrow habitat requirements for nesting and incubation. Whereas juveniles can be removed via electrofishing, angling, and perhaps minnow traps, options for targeting eggs and

larvae include the physical removal of eggs from nests, the chemical treatment of nests, nest predator management, sterilization, explosives (larvae only), water level management during nest guarding, and the removal of nest-guarding males. Although angling is itself an inefficient removal method (Boucher 2006; Gomez and Wilkinson 2008), angling may be an effective means of inducing nest failure. Generally, the aggression of nest-guarding males makes them easy to angle during spawning season (Ridgway et al. 1991). Both field and modeling studies involving *M. dolomieu* suggest that even short handling times during catch-and-release angling can increase the risk of nest predation (Ridgway and Shuter 1997; Suski et al. 2003; Steinhart et al. 2005), and that released bass can be too exhausted to adequately defend their nests (Kieffer et al. 1995; Hinch and Collins 1991; Hanson et al. 2008).

Conclusion

Human introductions and global climate change are facilitating the spread of *M. dolomieu* beyond their native range in Canada and the United States, and into other countries around the world. Invasive *M. dolomieu* can impact native food webs through predation, competition, and hybridization, and can even extirpate native fishes. These impacts are well documented and fairly well understood, but our ability to control invasive *M. dolomieu* is severely limited. Although numerous options for *M. dolomieu* control exist, few have been tested or developed and even fewer have been successful.

To improve *M. dolomieu* control, we recommend integrated pest management plans that include several nest failure strategies, perhaps in combination with other options (e.g., adult removal). For example, using catch-and-keep angling to remove nest-guarding males while simultaneously enhancing native nest predators is probably more effective at inducing nest failure than either of these options alone. Additionally, because a small subset of spawning males (~5%) can produce over half of a population's young of year (Gross and Kapuscinski 1997), we recommend using genetic techniques to identify and subsequently target the most productive males. We also recommend research to develop control methods that are not yet available for *M. dolomieu* (e.g., pathogens, parasites, pheromone traps). Of course, the combination of control options to use in an invaded system also depends on environmental constraints (e.g., lentic

vs. lotic, depth, structural complexity, substrate type, ecology), logistic constraints (e.g., budget, timeline, available infrastructure/equipment), and social factors (e.g., acceptability, political climate). *M. dolomieu* control may be unnecessary or impossible in some systems. For those systems in which control is an option, it is important to learn from previous attempts (*M. dolomieu* or other species) and practice adaptive management (Pine et al. 2009; Zipkin et al. 2009).

It is also important to maintain realistic expectations: *M. dolomieu* extirpation is unlikely except for in small and/or isolated systems, and *M. dolomieu* suppression will require multiple applications, probably in perpetuity (even rotenone can be <100% effective; Wydoski and Wiley 1999). For these reasons, in addition to sound research and management, we stress the importance of preventing *M. dolomieu* (re)introductions, for example through bait regulations and public awareness campaigns. Creative, comprehensive solutions need to be developed and implemented to minimize further spread of invasive *M. dolomieu* and mitigate their impacts on native ecosystems.

Acknowledgements

This work was funded by the National Science Foundation's Integrated Graduate Education and Research Traineeship (NSF-IGERT) program and the Conservation Biology graduate program of the University of Minnesota. We thank Ray Newman, Bruce Vondracek, Brian Shuter, Kyle Chezick, Justine Koch, and two anonymous reviewers for helpful comments on earlier versions of this manuscript. We also thank Jason Papenfuss for assistance with ArcGIS, Nick Phelps for insight on fish pathogens, and Fernanda Cabrini Araujo for her help searching the literature.

References

- Abrams PA (2009) When does greater mortality increase population size? The long history and diverse mechanisms underlying the hydra effect. *Ecology Letters* 12(5): 462–474, <http://dx.doi.org/10.1111/j.1461-0248.2009.01282.x>
- Allardi J, Keith P (1991) Atlas préliminaire des poissons d'eau douce de France. *Coll. Patrimoines Naturels* vol. 4 Secrétariat Faune Flore, Muséum national d'Histoire naturelle, Paris, 234 pp
- Andrews S (1985) Aquatic species introduced to Fiji. *Domodomo* 3(2): 67–82
- Archdeacon TP, Davenport SR (2010) Predation by age-0 smallmouth bass (*Micropterus dolomieu*) on bigscale logperch (*Percina macrolepida*) in the Pecos River, New Mexico. *Southwestern Naturalist* 55(1): 120–122, <http://dx.doi.org/10.1894/GG-40.1>
- Arnekleiv JV, Dolmen D, Ronning L (2001) Effects of rotenone treatment on mayfly drift and standing stocks in two Norwegian rivers. In: Dominguez E (ed), Trends in Research in Ephemeroptera and Plecoptera. Kluwer Academic/ Plenum Publishers, pp 77–88, http://dx.doi.org/10.1007/978-1-4615-1257-8_11

- Bacula TD, Blackwell BG, Willis DW (2011) Smallmouth bass seasonal dynamics in northeastern South Dakota glacial lakes. *Journal of Freshwater Ecology* 26(3): 345–356, <http://dx.doi.org/10.1080/02705060.2011.559744>
- Bajer PG, Chizinski CJ, Sorensen PW (2011) Using the Judas technique to locate and remove wintertime aggregations of invasive common carp. *Fisheries Management and Ecology* 18(6):497–505, <http://dx.doi.org/10.1111/j.1365-2400.2011.00805.x>
- Baker G, Darby N, Williams T, Wullschleger J (2008) Bonneville cutthroat trout restoration project—Great Basin National Park. Natural Resource Report NPS/NRPC/NRR–2008/055, US National Park Service, Fort Collins, Colorado
- Baxter II L, Hays EE, Hampson GR, Backus RH (1982) Mortality of fish subjected to explosive shock as applied to oil well severance on Georges Bank. Woods Hole Oceanographic Institute. Technical report No. WHOI-82-54, 73 pp
- Bayley PB, Austen DJ (1988) Comparison of detonating cord and rotenone for sampling fish in warmwater impoundments. *North American Journal of Fisheries Management* 8: 310–316, [http://dx.doi.org/10.1577/1548-8675\(1988\)008<0310:CODCAR>2.3.CO;2](http://dx.doi.org/10.1577/1548-8675(1988)008<0310:CODCAR>2.3.CO;2)
- Beamesderfer RC, Ward DL, Nigro AA (1996) Evaluation of the biological basis for a predator control program on northern squawfish (*Ptychocheilus oregonensis*) in the Columbia and Snake rivers. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 2898–2908, <http://dx.doi.org/10.1139/f96-225>
- Borden WC, Krebs RA (2009) Phylogeography and postglacial dispersal of smallmouth bass (*Micropterus dolomieu*) into the Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 66: 2142–2156, <http://dx.doi.org/10.1139/F09-155>
- Boucher DP (2005) Rapid River and Pond in the River Fisher Investigations. Report. Maine Department of Inland Fisheries and Wildlife. Fishery Progress Report Series No. 05-1, 49 pp
- Boucher DP (2006) A report on fish barrier needs and feasibility conducted on certain waters in the upper Androscoggin River drainage. Maine Department of Inland Fisheries and Wildlife, Rangeley Lakes Region. Unpublished report, 6 pp
- Bower S (1897) The propagation of small mouth black bass. *Transactions of the American Fisheries Society* 25: 127–136 [http://dx.doi.org/10.1577/1548-8659\(1896\)26\[127:TPOSMB\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1896)26[127:TPOSMB]2.0.CO;2),
- Brown CJD, Ball RC (1943) An experiment on the use of derris root (rotenone) on the fish and fish-food organisms of Third Sister Lake. *Transactions of the American Fisheries Society* 72: 267–284, [http://dx.doi.org/10.1577/1548-8659\(1942\)72\[267:AEITUO\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1942)72[267:AEITUO]2.0.CO;2)
- Brown TG, Runciman B, Pollard S, Grant ADA, Bradford MJ (2009) Biological synopsis of smallmouth bass (*Micropterus dolomieu*). Canadian Manuscript Report of Fisheries and Aquatic Sciences 2887: v + 50 pp
- Buhle ER, Margolis M, Ruesink JL (2005) Bang for buck: Cost-effective control of invasive species with different life histories. *Ecological Economics* 52(3): 355–366, <http://dx.doi.org/10.1016/j.ecolecon.2004.07.018>
- Burdick BD (2008) Removal of smallmouth bass and four other centrarchid fishes from the Upper Colorado and Lower Gunnison Rivers: 2004–2006. United States Fish and Wildlife Service: Upper Colorado River endangered fish recovery program. Project No. 126, 61 pp
- Cambray JA (2003) Impact on indigenous species biodiversity caused by the globalisation of alien recreational freshwater fisheries. *Hydrobiologia* 500(1–3):217–230, <http://dx.doi.org/10.1023/A:1024648719995>
- Carey MP, Sanderson BL, Friesen TA, Barnas KA, Olden JD (2011) Smallmouth bass in the Pacific Northwest: A threat to native species; a benefit for anglers. *Reviews in Fisheries Science* 19(3): 305–315, <http://dx.doi.org/10.1080/10641262.2011.598584>
- Chen RJ, Hunt KM, Ditton RB (2003) Estimating the economic impacts of a trophy largemouth bass fishery: Issues and applications. *North American Journal of Fisheries Management* 23(3): 835–844, <http://dx.doi.org/10.1577/M02-014>
- Chu C, Mandrak NE, Minns CK (2005) Potential impacts of climate change on the distributions of several common and rare freshwater fishes in Canada. *Diversity and Distributions* 11(4): 299–310, <http://dx.doi.org/10.1111/j.1366-9516.2005.00153.x>
- Crossman EJ (1991) Introduced freshwater fishes: A review of the North American perspective with emphasis on Canada. *Canadian Journal of Fisheries and Aquatic Sciences* 48(S1): 46–57, <http://dx.doi.org/10.1139/f91-303>
- Curry-Lindahl K (1966) Fiskarna i färg, 6th edn. Almqvist & Wiksell/Gerbers Förlag AB, Stockholm, Sweden, 242 pp
- Cushman OP (1917) Propagation of small-mouth black bass. *Transactions of the American Fisheries Society* 46(2): 113–116, [http://dx.doi.org/10.1577/1548-8659\(1916\)46\[113:POSB\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1916)46[113:POSB]2.0.CO;2)
- Davis HS (1937) A gill disease of the smallmouth blackbass. *The Progressive Fish-Culturist* 4(27): 7–11, [http://dx.doi.org/10.1577/1548-8640\(1937\)427\[7:AGDOTS\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8640(1937)427[7:AGDOTS]2.0.CO;2)
- Davis HS (1942) A suctorian parasite of the smallmouth black bass, with remarks on other suctorian parasites of fishes. *Transactions of the American Microscopical Society* 61(4): 309–327, <http://dx.doi.org/10.2307/3222897>
- Dawson VK, Kolar CS (eds) (2003) Integrated management techniques to control nonnative fishes. U.S. Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin, 146 pp, Appendixes A–F
- Dextrase A, Mandrak N (2006) Impacts of alien invasive species on freshwater fauna at risk in Canada. *Biological Invasions* 8(1): 13–24, <http://dx.doi.org/10.1007/s10530-005-0232-2>
- Dey CJ, O'Connor CM, Gilmour KM, Van Der Kraak G, Cooke SJ (2010) Behavioral and physiological responses of a wild teleost fish to cortisol and androgen manipulation during parental care. *Hormones and Behavior* 58: 599–605, <http://dx.doi.org/10.1016/j.yhbeh.2010.06.016>
- Dinger EC, Marks JC (2007) Effects of high levels of antimycin A on aquatic invertebrates in a warmwater Arizona stream. *North American Journal of Fisheries Management* 27: 1243–1256, <http://dx.doi.org/10.1577/M06-099.1>
- Drake DAR (2011) Quantifying the likelihood of human-mediated movements of species and pathogens: the baitfish pathway in Ontario as a model system. PhD Thesis, University of Toronto, Toronto, ON
- Dunlop ES, Orendorf JA, Shuter BJ, Rodd FH, Ridgway MS (2005a) Diet and divergence of introduced smallmouth bass (*Micropterus dolomieu*) populations. *Canadian Journal of Fisheries and Aquatic Sciences* 62(8): 1720–1732, <http://dx.doi.org/10.1139/f05-089>
- Dunlop ES, Shuter BJ, Ridgway MS (2005b) Isolating the influence of growth rate on maturation patterns in the smallmouth bass (*Micropterus dolomieu*). *Canadian Journal of Fisheries and Aquatic Sciences* 62(4): 844–853, <http://dx.doi.org/10.1139/f05-045>
- Earle JE, Lajeunesse BL (2007) Evaluation of a brook trout removal project to establish westslope cutthroat trout in Canmore Creek, Alberta. In Proceedings of Wild Trout IX symposium: Sustaining wild trout in a changing world, pp 9–12
- Edwards RJ (1980) The ecology and geographic variation of the Guadalupe bass, *Micropterus treculi*. PhD Thesis, University of Texas, Austin, TX
- Eiswerth ME, Johnson WS (2002) Managing nonindigenous invasive species: Insights from dynamic analysis. *Environmental and Resource Economics* 23(3): 319–342, <http://dx.doi.org/10.1023/A:1021275607224>
- FAO (1997) FAO Database on Introduced Aquatic Species, FAO, Rome

- Faulkner SG, Welz M, Tonn WM, Schmitt DR (2008) Effects of simulated blasting on mortality of rainbow trout eggs. *Transactions of the American Fisheries Society* 137: 1–12, <http://dx.doi.org/10.1577/T07-035.1>
- Fayram AH, Hansen MJ, Ehlinger TJ (2005) Interactions between walleyes and four fish species with implications for walleye stocking. *North American Journal of Fisheries Management* 25: 1321–1330, <http://dx.doi.org/10.1577/M04-203.1>
- Fermon Y (1997) Les Haplochromis spp. (Teleostei, Cichlidae) des zones rocheuses du Mwanza Gulf, lac Victoria, Tanzanie: Structure des communautés et écomorphologie. Thèse présentée pour l'obtention du Diplôme de Doctorat du Muséum National d'Histoire naturelle. Thèses et Documents Microfichés No. 157. ORSTOM, Paris
- Finlayson BJ, Schnick RA, Cailteux RL, Demong L, Horton WD, McClay W, Thompson CW, Tichacek GJ (2000) Rotenone use in fisheries management: Administrative and technical guidelines manual. American Fisheries Society, Bethesda, Maryland
- Finlayson BJ (2001) Introduction. In: Cailteux RL, Demong L, Finlayson BJ, Horton W, McClay W, Schnick RA, Thompson C (eds), Rotenone in fisheries: Are the rewards worth the risk? American Fisheries Society, Bethesda, Maryland, pp 1–3
- Finlayson BJ, Schnick RA, Cailteux RL, Demong L, Horton WD, McClay W, Thompson CW (2002) Assessment of Antimycin A use in fisheries and its potential for reregistration. *Fisheries* 27: 10–18, [http://dx.doi.org/10.1577/1548-8446\(2002\)027<0010:AOAAU>2.0.CO;2](http://dx.doi.org/10.1577/1548-8446(2002)027<0010:AOAAU>2.0.CO;2)
- Fricke R (1999) Fishes of the Mascarene Islands (Réunion, Mauritius, Rodriguez): An annotated checklist, with descriptions of new species. In: These Zoologicae, Volume 31. Koeltz Scientific Books, Koeningstein, 759 pp
- Fuller P, Cannister M (2011) *Micropterus dolomieu*. USGS Nonindigenous Aquatic Species Database, Gainesville, FL Available from: <http://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=396>
- Galster BJ, Wuellner MR, Graeb BD (2012) Walleye *Sander vitreus* and smallmouth bass *Micropterus dolomieu* interactions: An historic stable-isotope analysis approach. *Journal of Fish Biology* 81(1): 135–147, <http://dx.doi.org/10.1111/j.1095-8649.2012.03318.x>
- Gomez L, Wilkinson T (2008) A preliminary assessment of smallmouth bass in the beaver creek system. British Columbia Ministry of Environment, Cariboo Region
- Grizzle JM, Altinok I, Noyes AD (2003) PCR method for detection of largemouth bass virus. *Diseases of Aquatic Organisms* 54: 29–33, <http://dx.doi.org/10.3354/dao054029>
- Gross ML, Kapuscinski AR (1997) Reproductive success of smallmouth bass estimated and evaluated from family-specific DNA fingerprints. *Ecology* 78(5): 1424–1430, [http://dx.doi.org/10.1890/0012-9658\(1997\)078\[1424:RSOSBE\]2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(1997)078[1424:RSOSBE]2.0.CO;2)
- Halfyard EA (2010) A review of options for the containment, control and eradication of illegally introduced smallmouth bass (*Micropterus dolomieu*). Canadian Technical Report of Fisheries and Aquatic Sciences 2865, vi + 71 pp
- Hanel L (2003) The ichthyofauna of the Czech Republic: Development and present state. *Matthias Belvis University Proceedings* 3(1): 41–71
- Hanson KC, Gravel M, Redpath T, Cooke SJ, Siepker MJ (2008) Latitudinal variation in physiological and behavioral responses of nest-guarding smallmouth bass to common recreational angling practices. *Transactions of the American Fisheries Society* 137(5): 1558–1566, <http://dx.doi.org/10.1577/T07-228.1>
- Hawkins J, Walford C, Hill A (2008) Smallmouth bass control in the middle Yampa River, 2003–2007. United States Fish and Wildlife Service: Colorado State University, Fort Collins, Colorado to Upper Colorado River Endangered Fish Recovery Program. Final report, Larval Fish Laboratory Contribution 154, Denver, Colorado
- Hayes DB, Ferreri CP, Taylor WW (1996) Active fish capture methods. In: Murphy R, Willis DW (eds), *Fisheries Techniques*, 2nd edition. American Fisheries Society, Bethesda, Maryland, pp 193–220
- Heman ML, Campbell RS, Redmond LC (1969) Manipulation of fish populations through reservoir drawdown. *Transactions of the American Fisheries Society* 98: 293–304, [http://dx.doi.org/10.1577/1548-8659\(1969\)98\[293:MOFPTR\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1969)98[293:MOFPTR]2.0.CO;2)
- Henshall JA (1889) Book of the black bass. The Robert Clarke Company, Cincinnati, USA, 463 pp
- Henshall JA (1903) Bass, pike, perch and others. The Macmillan Company, New York, USA, 410 pp
- Hey S (1926) Preliminary Report on the Inland Waters of South Africa with Regard to the Suitability for the Introduction of Edible Fish. Department of Mines and Industries, Cape Town, South Africa, 140 pp
- Hinch SG, Collins NC (1991) Importance of diurnal and nocturnal nest defense in the energy budget of male smallmouth bass: Insights from direct video observations. *Transactions of the American Fisheries Society* 120(5): 657–663, [http://dx.doi.org/10.1577/1548-8659\(1991\)120<0657:IODANN>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1991)120<0657:IODANN>2.3.CO;2)
- Hodde MS (2004) Restoring balance: Using exotic species to control invasive exotic species. *Conservation Biology* 18: 38–49, <http://dx.doi.org/10.1111/j.1523-1739.2004.00249.x>
- Hunter LA (1988) Status of the endemic Atilán grebe of Guatemala: Is it extinct? *Condor* 90(4): 906–912, <http://dx.doi.org/10.2307/1368847>
- Iguchi KI, Yodo T, Matsubara N (2004a) Spawning and brood defense of smallmouth bass under the process of invasion into a novel habitat. *Environmental Biology of Fishes* 70(3): 219–225, <http://dx.doi.org/10.1023/B:EBFI.0000003337.44116.e8>
- Iguchi KI, Matsubara N, McNysset KM, Peterson AT, Scachetti-Pereira R, Powers KA, Vieglais DA, Wiley EO, Yodo T (2004b) Predicting invasions of North American basses in Japan using native range data and a genetic algorithm. *Transactions of the American Fisheries Society* 133(4): 845–854, <http://dx.doi.org/10.1577/T03-172.1>
- Iguchi KI, Yodo T (2004) Impact of indigenous egg eaters on the early survival of exotic smallmouth bass. *Ecological Research* 19(5): 469–474, <http://dx.doi.org/10.1111/j.1440-1703.2004.00660.x>
- Inskip PD, Magnuson JG (1983) Changes in fish populations over an 80-year period: Big Pine Lake, Wisconsin. *Transactions of the American Fisheries Society* 112: 378–389, [http://dx.doi.org/10.1577/1548-8659\(1983\)112<378:CIFPOA>2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1983)112<378:CIFPOA>2.0.CO;2)
- Jackson DA, Peres-Neto P, Olden JD (2001) What controls who is where in freshwater fish communities — The roles of biotic, abiotic, and spatial factors. *Canadian Journal of Fisheries and Aquatic Sciences* 58(1): 157–170
- Jackson DA (2002) Ecological effects of *Micropterus* introductions: The dark side of black bass. *American Fisheries Society Symposium* 31: 221–232
- Jackson DA, Mandrak NE (2002) Changing fish biodiversity: Predicting the loss of cyprinid biodiversity due to global climate change. *American Fisheries Society Symposium* 32: 89–98
- Johnson BM, Martinez PJ, Hawkins JA, Bestgen KR (2008) Ranking predatory threats by nonnative fishes in the Yampa River, Colorado, via bioenergetics modeling. *North American Journal of Fisheries Management* 28(6): 1941–1953, <http://dx.doi.org/10.1577/M07-199.1>
- Johnson FH, Hale JG (1977) Interrelations between walleye (*Stizostedion vitreum vitreum*) and smallmouth bass (*Micropterus dolomieu*) in four northeastern Minnesota lakes, 1948–69. *Journal of the Fisheries Research Board of Canada* 34: 1626–1632, <http://dx.doi.org/10.1139/f77-227>

- Keevin TM, Hempen GL, Davinroy RD, Rapp RJ (2002) The use of high explosives to conduct a fisheries survey at a bendway weir field on the middle Mississippi River. In: Proceedings of the Annual General Meeting of the International Society of Explosives Engineers, pp 380–391
- Kempinger JJ, Carline RF (1977) Dynamics of the walleye (*Stizostedion vitreum vitreum*) population in Escanaba Lake, Wisconsin, 1955–72. *Journal of the Fisheries Research Board of Canada* 34: 1800–1811, <http://dx.doi.org/10.1139/f77-246>
- Kieffer JD, Kubacki MR, Phelan FJS, Philipp DP, Tufts BL (1995) Effects of catch-and-release angling on nesting male smallmouth bass. *Transactions of the American Fisheries Society* 124(1): 70–76, [http://dx.doi.org/10.1577/1548-8659\(1995\)124<0070:EOCARA>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1995)124<0070:EOCARA>2.3.CO;2)
- Kiesecker JM, Blaustein AR (2008) Effects of introduced bullfrogs and smallmouth bass on microhabitat use, growth, and survival of native red-legged frogs (*Rana aurora*). *Conservation Biology* 12(4): 776–787, <http://dx.doi.org/10.1111/j.1523-1739.1998.97125.x>
- Kitazima J, Mori S (2011) Re-introduction and re-enforcement of oily bitterling in conjunction with local communities in Northern Mie, Japan. In: Soorae PS (ed), Global re-introduction perspectives: 2011. More case studies from around the globe. IUCN/SSC Re-introduction Specialist Group and Environment Agency-Abu Dhabi, Gland, Switzerland and Abu Dhabi, UAE: pp 65–70
- Kleinschmidt Energy and Water Resource Consultants (2008) Smallmouth bass / brook trout habitat manipulation studies in Rapid River TWP C and Upton, Oxford County, Maine - 2007 progress report. Submitted to Rapid River Coalition, Maine Department of Inland Fisheries and Wildlife and Trout Unlimited, 39 pp
- Kolar CS, Chapman DC, Courtenay WR, Housel CM, Williams JD, Jennings DP (2005) Asian carps of the genus *Hypophthalmichthys* (Pisces, Cyprinidae) – A biological synopsis and environmental risk assessment. Report to the U.S. Fish and Wildlife Service, 183 pp
- Kullander SO, Nyman L, Jilg K, Delling B (2012) Nationalnyckeln till Sveriges flora och fauna. Strålfeniga fiskar. Actinopterygii. ArtDatabanken, SLU, Uppsala, 517 pp
- Kulp MA, Moore SE (2000) Multiple electrofishing removals for eliminating rainbow trout in a small southern Appalachian stream. *North American Journal of Fisheries Management* 20: 259–266, [http://dx.doi.org/10.1577/1548-8675\(2000\)020<0259:MERFER>2.0.CO;2](http://dx.doi.org/10.1577/1548-8675(2000)020<0259:MERFER>2.0.CO;2)
- Kuronuma K (1961) A check list of fishes of Vietnam. Division of Agriculture and Natural Resources, United States Operations Mission to Vietnam. United States Consultants, Inc.; International Cooperation Administration Contract-IV-153, 66 pp
- Lee DP (2001) Northern Pike Control in Lake Davis, California. In: Cailteux RL, Demong L, Finlayson BJ, Horton W, McClay W, Schnick RA, Thompson C (eds), Rotenone in fisheries: Are the rewards worth the risk? American Fisheries Society, Bethesda, Maryland, pp 55–62
- Lennon RE, Hunn JB, Schnick RA, Burrell RM (1971) Reclamation of lakes, ponds and streams with fish toxicants: A review. Food and Agriculture Organization of the United Nations (FAO). Fisheries Technical Paper No. 100, 99 pp
- Liao H, Pierce CL, Larscheid JG (2004) Consumption dynamics of the adult piscivorous fish community in Spirit Lake, Iowa. *North American Journal of Fisheries Management* 24: 890–902, <http://dx.doi.org/10.1577/M02-178.1>
- Ling N (2001) Rotenone: a review of its toxicity and use for fisheries management. New Zealand Department of Conservation. Science for Conservation, Report No. 211, Wellington, New Zealand, 39 pp
- Litvak MK, Mandrak NE (1993) Ecology of freshwater baitfish use in Canada and the United States. *Fisheries* 18(12): 6–13, [http://dx.doi.org/10.1577/1548-8446\(1993\)018<0006:EOFBUI>2.0.CO;2](http://dx.doi.org/10.1577/1548-8446(1993)018<0006:EOFBUI>2.0.CO;2)
- Lotufo GR, Lydy MJ (2005) Comparative toxicokinetics of explosive compounds in sheepshead minnows. *Archives of Environmental Contamination and Toxicology* 49: 206–214, <http://dx.doi.org/10.1007/s00244-004-0197-7>
- Lydell D (1902) The habits and culture of the black bass. *Transactions of the American Fisheries Society* 31(1): 45–73, [http://dx.doi.org/10.1577/1548-8659\(1902\)32\[45:THACOT\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1902)32[45:THACOT]2.0.CO;2)
- Lyons J (2011) Smallmouth bass, *Micropterus dolomieu*. Fishes of Wisconsin E-book. Wisconsin Department of Natural Resources, Madison, and U.S. Geological Survey, Middleton, WI. <http://www.fow-ebook.us> (Accessed 21 February 2012)
- MacRae PSD, Jackson DA (2001) The influence of smallmouth bass (*Micropterus dolomieu*) predation and habitat complexity on the structure of littoral zone fish assemblages. *Canadian Journal of Fisheries and Aquatic Sciences* 58(2): 342–351
- Maezono Y, Miyashita T (2003) Community-level impacts induced by introduced largemouth bass and bluegill in farm ponds in Japan. *Biological Conservation* 109(1): 111–121, [http://dx.doi.org/10.1016/S0006-3207\(02\)00144-1](http://dx.doi.org/10.1016/S0006-3207(02)00144-1)
- Maezono Y, Kobayashi R, Kusahara M, Miyashita T (2005) Direct and indirect effects of exotic bass and bluegill on exotic and native organisms in farm ponds. *Ecological Applications* 15: 638–650, <http://dx.doi.org/10.1890/02-5386>
- Marcinkevage Miller AC (2007) An individual-based model of fish habitat selection for network-scale watershed assessment. ProQuest Information and Learning Company, Ann Arbor, MI
- Masuda H, Amaoka K, Araga C, Uyeno T, Yoshino T (1984) The fishes of the Japanese Archipelago, Volume 1. Tokai University Press, Tokyo, Japan, 437 pp
- McClay W (2000) Rotenone use in North America (1988–1997). *Fisheries* 5: 5–21
- McCull KA, Sunarto A, Williams LM, Crane MSJ (2007) Koi herpes virus: Dreaded pathogen or white knight? *Aquaculture Health International* 9: 4–6
- McCormick JH, Stokes GN (1982) Intraovarian invasion of smallmouth bass oocytes by *Proteocephalus ambloplitis* (Cestoda). *Journal of Parasitology* 68(5): 975–976, <http://dx.doi.org/10.2307/3281024>
- Meronek TG, Bouchard PM, Buckner ER, Burri TM, Demmerly KK, Hatleli DC, Klumb RA, Schmidt SH, Coble DW (1996) A review of fish control projects. *North American Journal of Fisheries Management* 16: 63–74, [http://dx.doi.org/10.1577/1548-8675\(1996\)016<0063:AROFPC>2.3.CO;2](http://dx.doi.org/10.1577/1548-8675(1996)016<0063:AROFPC>2.3.CO;2)
- Metzger RJ, Shafland PL (1986) Use of detonating cord for sampling fish. *North American Journal of Fisheries Management* 6(1): 113–118, [http://dx.doi.org/10.1577/1548-8659\(1986\)6<113:UODCF>2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1986)6<113:UODCF>2.0.CO;2)
- Moore SE, Larson GL, Ridley B (1986) Population control of exotic rainbow in streams of a natural area park. *Environmental Management* 10: 215–219, <http://dx.doi.org/10.1007/BF01867359>
- Moore SE, Kulp MA, Hammonds J, Rosenlund B (2005) Restoration of Sams Creek and an assessment of brook trout restoration methods – Great Smoky Mountains National Park. US National Park Service. Technical Report NPS/NRWRD/NRTR-2005/342
- Morbey YE, Vascotto K, Shuter BJ (2007) Dynamics of piscivory by lake trout following a smallmouth bass invasion: A historical reconstruction. *Transactions of the American Fisheries Society* 136(2): 477–483, <http://dx.doi.org/10.1577/T06-070.1>

- Morrison BRS (1979) An investigation into the effects of the piscicide antimycin A on the fish and invertebrates of a Scottish stream. *Fisheries Management* 10: 111–122
- Moyle PB (1976) Fish introductions in California: History and impact on native fishes. *Biological Conservation* 9: 101–118, [http://dx.doi.org/10.1016/0006-3207\(76\)90043-4](http://dx.doi.org/10.1016/0006-3207(76)90043-4)
- Mukai T, Tsukahara K, Miwa Y (2011) The re-introduction of the Ushimotsugo minnow in Gifu Prefecture, Japan. In: Soorae PS (ed), Global re-introduction perspectives: 2011. More case studies from around the globe. IUCN/SSC Re-introduction Specialist Group and Environment Agency-Abu Dhabi, Gland, Switzerland and Abu Dhabi, UAE, pp 54–58
- Munther GL (1970) Movement and distribution of smallmouth bass in the Middle Snake River. *Transactions of the American Fisheries Society* 99(1): 44–53, [http://dx.doi.org/10.1577/15488659\(1970\)99<44:MADOSB>2.0.CO;2](http://dx.doi.org/10.1577/15488659(1970)99<44:MADOSB>2.0.CO;2)
- Page LM, Burr BM (1991) A field guide to freshwater fishes of North America north of Mexico. Houghton Mifflin Company, Boston, MA, 432 pp
- Peterson JT, Kwak TJ (1999) Modeling the effects of land use and climate change on riverine smallmouth bass. *Ecological Applications* 9(4): 1391–1404, [http://dx.doi.org/10.1890/1051-0761\(1999\)009\[1391:MTEOLUJ\]2.0.CO;2](http://dx.doi.org/10.1890/1051-0761(1999)009[1391:MTEOLUJ]2.0.CO;2)
- Pine WE, Martell SJD, Walters CJ, Kitchell JF (2009) Counterintuitive responses of fish populations to management actions: Some common causes and implications for predictions based on ecosystem modeling. *Fisheries* 34(4): 165–180, <http://dx.doi.org/10.1577/1548-8446-34.4.165>
- Rahel FJ (2000) Homogenization of fish faunas across the United States. *Science* 288: 854–856, <http://dx.doi.org/10.1126/science.288.5467.854>
- Rahel FJ (2004) Unauthorized fish introductions: Fisheries management of the people, for the people, or by the people? *American Fisheries Society Symposium* 44: 431–433
- Rahel FJ, Olden JD (2008) Assessing the effects of climate change on aquatic invasive species. *Conservation Biology* 22: 521–533, <http://dx.doi.org/10.1111/j.1523-1739.2008.00950.x>
- Rawson DS (1945) The experimental introduction of smallmouth black bass into lakes of the Prince Albert National Park, Saskatchewan. *Transactions of the American Fisheries Society* 73(1): 19–31, [http://dx.doi.org/10.1577/1548-8659\(1943\)73\[19:TEIOSB\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1943)73[19:TEIOSB]2.0.CO;2)
- Ridgway MS, Shuter BJ, Post EE (1991) The relative influence of body size and territorial behaviour on nesting asynchrony in male smallmouth bass, *Micropterus dolomieu* (Pisces: Centrarchidae). *Journal of Animal Ecology* 60(2): 665–681, <http://dx.doi.org/10.2307/5304>
- Ridgway MS, Shuter BJ (1997) Predicting the effects of angling for nesting male smallmouth bass on production of age-0 fish with an individual-based model. *North American Journal of Fisheries Management* 17(2): 568–580, [http://dx.doi.org/10.1577/1548-8675\(1997\)017<0568:PTEOAF>2.3.CO;2](http://dx.doi.org/10.1577/1548-8675(1997)017<0568:PTEOAF>2.3.CO;2)
- Ridgway MS, Shuter BJ, Middel TA, Gross ML (2002) Spatial ecology and density-dependent processes in smallmouth bass: The juvenile transition hypothesis. *American Fisheries Society Symposium* 31: 47–60
- Rieman BE, Beamesderfer RC, Vigg S, Poe TP (1991) Estimated loss of juvenile salmonids to predation by northern squawfish, walleyes, and smallmouth bass in John Day Reservoir, Columbia River. *Transactions of the American Fisheries Society* 120(4): 448–458, [http://dx.doi.org/10.1577/1548-8659\(1991\)120<0448:ELOJST>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1991)120<0448:ELOJST>2.3.CO;2)
- Rinne JN (2001) Nonnative, predatory fish removal and native fish response, upper Verde River, Arizona: Preliminary results. *Hydrology and Water Resources in Arizona and the Southwest* 31: 29–36
- Ripple R (1908) Experiments in rearing small-mouth black bass. *Transactions of the American Fisheries Society* 37(1): 126–131, [http://dx.doi.org/10.1577/1548-8659\(1908\)38\[126:EIRSB\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1908)38[126:EIRSB]2.0.CO;2)
- Robillard MM, Fox MG (2006) Historical changes in abundance and community structure of warmwater piscivore communities associated with water clarity, nutrients, and temperature. *Canadian Journal of Fisheries and Aquatic Sciences* 63: 798–809, <http://dx.doi.org/10.1139/f05-259>
- Rogers KB, Bergersen EP (1995) Effects of a fall drawdown on movement of adult northern pike and largemouth bass. *North American Journal of Fisheries Management* 15: 596–600, [http://dx.doi.org/10.1577/1548-8675\(1995\)015<0596:EOAFDO>2.3.CO;2](http://dx.doi.org/10.1577/1548-8675(1995)015<0596:EOAFDO>2.3.CO;2)
- Rose KA (2005) Lack of relationship between simulated fish population responses and their life history traits: Inadequate models, incorrect analysis, or site-specific factors? *Canadian Journal of Fisheries and Aquatic Sciences* 62(4): 886–902, <http://dx.doi.org/10.1139/f05-049>
- Sanderson BL, Barnas KA, Wargo Rub AM (2009) Nonindigenous species of the Pacific Northwest: An overlooked risk to endangered salmon? *BioScience* 59(3): 245–256, <http://dx.doi.org/10.1525/bio.2009.59.3.9>
- Schlosser IJ (1987) The role of predation in age- and size-related habitat use by stream fishes. *Ecology* 68(3): 651–659, <http://dx.doi.org/10.2307/1938470>
- Schramm HL, Armstrong ML, Funicelli NA, Green DM, Lee DP, Manns RE, Taubert BD, Waters SJ (1991) The status of competitive sport fishing in North America. *Fisheries* 16(3): 4–12, [http://dx.doi.org/10.1577/1548-8446\(1991\)016<0004:TSCSFS>2.0.CO;2](http://dx.doi.org/10.1577/1548-8446(1991)016<0004:TSCSFS>2.0.CO;2)
- Scott WB, Crossman EJ (1973) Freshwater fishes of Canada. Bulletin of the Fisheries Research Board of Canada No. 184
- Settle LR, Govoni JJ, Greene MD, West MA (2002) The effects of underwater explosions on larval fish with implications for the Wilmington Harbor Project. U.S. Army Corps of Engineers. Wilmington, NC
- Sharma S, Jackson DA, Minns CK, Shuter BJ (2007) Will northern fish populations be in hot water because of climate change? *Global Change Biology* 13(10): 2052–2064, <http://dx.doi.org/10.1111/j.1365-2486.2007.01426.x>
- Sharma S, Herborg L, Theriault TW (2009a) Predicting introduction, establishment and potential impacts of smallmouth bass. *Diversity and Distributions* 15: 831–840, <http://dx.doi.org/10.1111/j.1472-4642.2009.00585.x>
- Sharma S, Jackson DA, Minns CK (2009b) Quantifying the potential effects of climate change and the invasion of smallmouth bass on native lake trout populations across Canadian lakes. *Ecography* 32(3): 517–525, <http://dx.doi.org/10.1111/j.1600-0587.2008.05544.x>
- Shroyer SM (2007) Induced winterkill as a management tool for reclaiming Minnesota walleye rearing ponds. Minnesota Department of Natural Resources. Investigational Report 547, St. Paul, Minnesota
- Shuter B J, Maclean JA, Fry FEJ, Regier HA (1980) Stochastic simulation of temperature effects on 1st-year survival of smallmouth bass. *Transactions of the American Fisheries Society* 109(1): 1–34, [http://dx.doi.org/10.1577/1548-8659\(1980\)109<1:SSOTEO>2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1980)109<1:SSOTEO>2.0.CO;2)
- Shuter BJ, Ihssen PE, Wales DL, Snucins EJ (1989) The effects of temperature, pH and water hardness on winter starvation of young-of-the-year smallmouth bass, *Micropterus dolomieu lacepede*. *Journal of Fish Biology* 35(6): 765–780, <http://dx.doi.org/10.1111/j.1095-8649.1989.tb03028.x>
- Smale MA, Rabeni CF (1995) Hypoxia and hyperthermia tolerances of headwater stream fishes. *Transactions of the American Fisheries Society* 124(5): 698–710, [http://dx.doi.org/10.1577/1548-8659\(1995\)124<0698:HAHTOH>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1995)124<0698:HAHTOH>2.3.CO;2)
- Smith MW (1940) Copper sulfate and rotenone as fish poisons. *Transactions of the American Fisheries Society* 69: 141–157, [http://dx.doi.org/10.1577/1548-8659\(1939\)69\[141:CSARAF\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1939)69[141:CSARAF]2.0.CO;2)
- Smith MW (1941) Treatment of Potter's Lake, New Brunswick, with rotenone. *Transactions of the American Fisheries Society* 70(1): 347–355, [http://dx.doi.org/10.1577/1548-8659\(1940\)70\[347:TOPLNB\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(1940)70[347:TOPLNB]2.0.CO;2)

- Sorensen PW, Stacey NE (2004) Brief review of fish pheromones and discussion of their possible uses in the control of non-indigenous teleost fishes. *N.Z. Journal of Marine and Freshwater Research* 38(3): 399–417, <http://dx.doi.org/10.1080/00288330.2004.9517248>
- Steinhart GB, Marschall EA, Stein RA (2004) Round goby predation on smallmouth bass offspring in nests during simulated catch-and-release angling. *Transactions of the American Fisheries Society* 133(1): 121–131, <http://dx.doi.org/10.1577/T03-020>
- Steinhart GB, Leonard NJ, Stein RA, Marschall EA (2005) Effects of storms, angling, and nest predation during angling on smallmouth bass (*Micropterus dolomieu*) nest success. *Canadian Journal of Fisheries and Aquatic Sciences* 62(11): 2649–2660, <http://dx.doi.org/10.1139/f05-171>
- Stevens CMJ, Bonsall MB (2011) The impact of alternative harvesting strategies in a resource-consumer metapopulation. *Journal of Applied Ecology* 48(1): 102–111, <http://dx.doi.org/10.1111/j.1365-2664.2010.01907.x>
- Suski CD, Svec JH, Ludden JB, Phelan FJS, Philipp DP (2003) The effect of catch-and-release angling on the parental care behavior of male smallmouth bass. *Transactions of the American Fisheries Society* 132(2): 210–218, [http://dx.doi.org/10.1577/1548-8659\(2003\)132<0210:TEOCAR>2.0.CO;2](http://dx.doi.org/10.1577/1548-8659(2003)132<0210:TEOCAR>2.0.CO;2)
- Taylor D, Katahira L (1988) Radio telemetry as an aid in eradicating remnant feral goats. *Wildlife Society Bulletin* 16: 297–299
- Teuscher D, Luecke C (1996) Competition between kokanees and Utah chub in Flaming Gorge Reservoir, Utah-Wyoming. *Transactions of the American Fisheries Society* 125(4): 505–511, [http://dx.doi.org/10.1577/1548-8659\(1996\)125<0505:CBKAUC>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1996)125<0505:CBKAUC>2.3.CO;2)
- Thresher RE (2008) Autocidal technology for the control of invasive fish. *Fisheries* 33: 113–121, <http://dx.doi.org/10.1577/1548-8446-33.3.114>
- Trumpickas J, Mandrak NE, Ricciardi A (2011) Nearshore fish assemblages associated with introduced predatory fishes in lakes. *Aquatic Conservation: Marine and Freshwater Ecosystems* 21: 338–347, <http://dx.doi.org/10.1002/aqc.1192>
- Twohey MB, Heinrich JW, Seelye JG, Fredricks KT, Bergstedt RA, Kaye CA, Scholefield RJ, McDonald RB, Christie GC (2003) The sterile-male-release technique in Great Lakes sea lamprey management. *Journal of Great Lakes Research* 29: 410–423, [http://dx.doi.org/10.1016/S0380-1330\(03\)70504-8](http://dx.doi.org/10.1016/S0380-1330(03)70504-8)
- Tyus HM, Saunders JF (2000) Nonnative fish control and endangered fish recovery: Lessons from the Colorado River. *Fisheries* 25(9): 17–24, [http://dx.doi.org/10.1577/1548-8446\(2000\)025<0017:NFCAEF>2.0.CO;2](http://dx.doi.org/10.1577/1548-8446(2000)025<0017:NFCAEF>2.0.CO;2)
- Vander Zanden MJ, Casselman JM, Rasmussen JB (1999) Stable isotope evidence for the food web consequences of species invasions in lakes. *Nature* 401: 464–467, <http://dx.doi.org/10.1038/46762>
- Vander Zanden MJ, Olden JD, Thorne JH, Mandrak NE (2004) Predicting occurrences and impacts of smallmouth bass introductions in north temperate lakes. *Ecological Applications* 14: 132–148, <http://dx.doi.org/10.1890/02-5036>
- Van Winkle W, Richter TJ, Chandler JA (2001) Relative potential consequences of alternative operational scenarios for centrarchid populations in Brownlee Reservoir. Idaho Power Company. Hells Canyon Complex Hydroelectric Project, Technical Report, Appendix E.3.1-5, Chapter 4, Boise, Idaho
- Verrill DD, Berry CR (1995) Effectiveness of an electrical barrier and lake drawdown for reducing common carp and bigmouth buffalo abundances. *North American Journal of Fisheries Management* 15: 137–141, [http://dx.doi.org/10.1577/1548-8675\(1995\)015<0137:EOAEB>2.3.CO;2](http://dx.doi.org/10.1577/1548-8675(1995)015<0137:EOAEB>2.3.CO;2)
- Wagner CM, Jones ML, Twohey MB, Sorensen PW (2006) A field test verifies that pheromones can be useful for sea lamprey (*Petromyzon marinus*) control in the Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 63(3): 475–479, <http://dx.doi.org/10.1139/f06-008>
- Ward D (2005) Selective removal of nonnative fishes using Supavern®: Toxicity screening for a candidate species-specific piscicide. *Journal of Freshwater Ecology* 20(4): 787–789, <http://dx.doi.org/10.1080/02705060.2005.9664807>
- Warner K (2005) Smallmouth bass introductions in Maine: History and management implications. *Fisheries* 30(11): 20–26, [http://dx.doi.org/10.1577/1548-8446\(2005\)30\[20:SBIM\]2.0.CO;2](http://dx.doi.org/10.1577/1548-8446(2005)30[20:SBIM]2.0.CO;2)
- Weidel BC, Josephson DC, Kraft CE (2007) Littoral fish community response to smallmouth bass removal from an Adirondack lake. *Transactions of the American Fisheries Society* 136: 778–789, <http://dx.doi.org/10.1577/T06-091.1>
- Welcomme RL (1988) International introductions of inland aquatic species. FAO Fisheries Technical Paper 294, 318 pp
- Whitmore DH (1983) Introgressive hybridization of smallmouth bass (*Micropterus dolomieu*) and Guadalupe bass (*Micropterus treculi*). *Copeia* (3): 672–679, <http://dx.doi.org/10.2307/1444331>
- Whitmore DH, Hellier TR (1988) Natural hybridization between largemouth and smallmouth bass (*Micropterus*). *Copeia* 1988(2): 493–496, <http://dx.doi.org/10.2307/1445895>
- Wright J (2000) Seasonal variation in trap net catches of smallmouth bass in two northwestern Ontario lakes. Ontario Ministry of Natural Resources. Northwest Science and Information, Aquatics Update, 1(4)
- Wydoski RS, Wiley RW (1999) Management of undesirable fish species. In: Kohler CC, Hubert WA (eds), Inland fisheries management in North America, 2nd edition. American Fisheries Society, Bethesda, Maryland, pp 403–430
- Yule DL, Luecke C (1993) Lake trout consumption and recent changes in the fish assemblage of Flaming Gorge Reservoir. *Transactions of the American Fisheries Society* 122: 1058–1069, [http://dx.doi.org/10.1577/1548-8659\(1993\)122<1058:LTARC>2.3.CO;2](http://dx.doi.org/10.1577/1548-8659(1993)122<1058:LTARC>2.3.CO;2)
- Zipkin EF, Sullivan PJ, Cooch EG, Kraft CE, Shuter BJ, Weidel BC (2008) Overcompensatory response of a smallmouth bass (*Micropterus dolomieu*) population to harvest: Release from competition? *Canadian Journal of Fisheries and Aquatic Sciences* 65(10): 2279–2292, <http://dx.doi.org/10.1139/F08-133>
- Zipkin EF, Kraft CE, Cooch EG, Sullivan PJ (2009) When can efforts to control nuisance and invasive species backfire? *Ecological Applications* 19(6): 1585–1595, <http://dx.doi.org/10.1890/08-1467.1>

Impacts of invasive smallmouth bass. Invasive *M. dolomieu* can disrupt the native ecology of the systems to which they have been introduced. Bass are voracious predators that can decrease the abundance of, change the habitat used by, and even extirpate small prey fish such as brook stickleback (*Culaea inconstans* Kirtland, 1840), fathead minnow (*Pimephales promelas* Rafinesque, 1820), pearl dace (*Margariscus margarita* Cope, 1867), finescale dace (*Phoxinus neogaeus* Cope, 1867) and northern redbelly dace (*Phoxinus eos* Cope, 1861) (MacRae. Table 2. Invasive fish control methods, examples of their use for smallmouth bass control, the bass life stage they target, and their pros and cons. This table draws from and builds upon a review by Halfyard (2010). Invasive smallmouth bass (*Micropterus dolomieu*): History, impacts, and control. Grace L. Loppnow, Kris Vascotto, Paul A. Venturelli. 2013. Ecological effects of *Micropterus* introductions: the dark side of black bass. In *Black Bass: Ecology, Conservation, and Management* (Phillip. D. A. Jackson. American Fisheries Society.